COUNTING ON YOU

Keep Children and Teens Healthy by Preventing
Multisystem Inflammatory Syndrome in Children (MIS-C)



Vaccinate to Prevent COVID-19 and MIS-C

If you think your child is showing symptoms of MIS-C, trust your instincts and contact your child's doctor, nurse, or clinic right away.

Children, adolescents, or young adults who develop certain symptoms after having COVID-19 might have MIS-C. They should see a doctor if they had COVID-19, or have been in close contact with someone who had COVID-19, within the past 6 weeks and now have symptoms.

Not all children will have the same symptoms.

It could be MIS-C if your child has an ongoing fever AND more than one of the following symptoms:

- Stomach Pain
- Bloodshot Eyes
- Vomiting and or Diarrhea
- Dizziness and or Lightheadedness
- Skin Rash





Let a healthcare provider know if your child has a fever and other symptoms; and has been in contact with the COVID-19 virus.

COUNTING ON YOU

Keep Children and Teens Healthy by Preventing Multisystem Inflammatory Syndrome in Children (MIS-C)



Vaccinate to Prevent COVID-19 and MIS-C

If you think your child is showing symptoms of MIS-C, trust your instincts and contact your child's doctor, nurse, or clinic right away.

Children, adolescents, or young adults who develop certain symptoms after having COVID-19 might have MIS-C. They should see a doctor if they had COVID-19, or have been in close contact with someone who had COVID-19, within the past 6 weeks and now have symptoms.

Not all children will have the same symptoms.

It could be MIS-C if your child has an ongoing fever AND more than one of the following symptoms:

- Stomach Pain
- Bloodshot Eyes
- Vomiting and or Diarrhea
- Dizziness and or Lightheadedness
- Skin Rash



MIS-C Fact Sheet (pa.gov



PA.GOV/COVID 1-877-PA-HEALTH (1-877-724-3258)

Let a healthcare provider know if your child has a fever and other symptoms; and has been in contact with the COVID-19 virus.

COUNTING ON YOU

Keep Children and Teens Healthy by Preventing Multisystem Inflammatory Syndrome in Children (MIS-C)



Vaccinate to Prevent COVID-19 and MIS-C

If you think your child is showing symptoms of MIS-C, trust your instincts and contact your child's doctor, nurse, or clinic right away.

Children, adolescents, or young adults who develop certain symptoms after having COVID-19 might have MIS-C. They should see a doctor if they had COVID-19, or have been in close contact with someone who had COVID-19, within the past 6 weeks and now have symptoms.

Not all children will have the same symptoms.

It could be MIS-C if your child has an ongoing fever AND more than one of the following symptoms:

- Stomach Pain
- Bloodshot Eyes
- Vomiting and or Diarrhea
- Dizziness and or Lightheadedness
- Skin Rash





Let a healthcare provider know if your child has a fever and other symptoms; and has been in contact with the COVID-19 virus.